

VYVGART Hytrulo®  
(efgartigimod alfa and hyaluronidase-qvfc)

VYVGART®  
(efgartigimod alfa-fcab)

Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Start the conversation with your doctor

**MY current MG-ADL score is:**  
(0 to 24)

**Don't know your MG-ADL score?**

Visit [MySymptomScore.com](https://mysymptomscore.com) to use an interactive tool to measure 8 signs or symptoms that are commonly affected in gMG.



**What goal would I choose now if MY gMG symptoms and daily abilities could be improved?**

---

---

**This goal is important to me because:**

---

---

**This is a goal I want to work on now because:**

---

---

**What are some other things I would think about doing in the future if my gMG symptoms and daily abilities could be improved?**

---

---

**Why do I want to work on these goals in the future?**

---

---



**Now that you've set your Personal Ability Goal, it's time to share it with your doctor**

**Important tip!** Your Personal Ability Goal may change over time. Make sure to set new goals and talk to your doctor about them.

This information has been provided for educational purposes only and is not meant as a substitute for professional medical advice. Remember to let your healthcare provider know if there are any changes to your MG-ADL score. AChR=acetylcholine receptor; MG-ADL=Myasthenia Gravis Activities of Daily Living.

## **What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?**

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.

It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

### **SELECT IMPORTANT SAFETY INFORMATION**

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

**Please see Important Safety Information on the next page. Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.**

## IMPORTANT SAFETY INFORMATION

Do not take VYVGART if you are allergic to efgartigimod alfa or any of the ingredients in VYVGART. Do not take VYVGART HYTRULO if you are allergic to efgartigimod alfa, hyaluronidase, or any of the ingredients in VYVGART HYTRULO. VYVGART or VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

### Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection or fever.
- have recently received or are scheduled to receive any vaccinations.
- have any history of allergic reactions.
- have kidney (renal) problems.
- are pregnant or plan to become pregnant. It is not known whether VYVGART or VYVGART HYTRULO will harm your unborn baby.
  - Pregnancy Exposure Registry. There is a pregnancy exposure registry for women who use VYVGART or VYVGART HYTRULO during pregnancy. The purpose of this registry is to collect information about your health and your baby. Your healthcare provider can enroll you in this registry. You may also enroll yourself or get more information about the registry by calling 1-855-272-6524 or going to [VYVGARTPregnancy.com](http://VYVGARTPregnancy.com)
- are breastfeeding or plan to breastfeed. It is not known if VYVGART or VYVGART HYTRULO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

### VYVGART or VYVGART HYTRULO can cause side effects which can be serious, including:

- **Infection.** VYVGART or VYVGART HYTRULO may increase the risk of infection. If you have an active infection, your healthcare provider should delay your treatment with VYVGART or VYVGART HYTRULO until your infection is gone. Tell your healthcare provider right away if you get any of the following signs and symptoms of an infection:
  - fever
  - chills
  - frequent and painful urination
  - cough
  - pain and blockage of nasal passages
  - wheezing
  - shortness of breath
  - sore throat
  - excess phlegm
  - nasal discharge

- **Allergic reactions (hypersensitivity reactions).** VYVGART or VYVGART HYTRULO can cause allergic reactions that can be severe. These reactions can happen during, shortly after, or weeks after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction with VYVGART or VYVGART HYTRULO:
  - rash
  - swelling of the face, lips, throat, or tongue
  - shortness of breath
  - trouble breathing
  - low blood pressure
  - fainting

- rash
- swelling of the face, lips, throat, or tongue
- shortness of breath
- trouble breathing
- low blood pressure
- fainting

An additional symptom of an allergic reaction with VYVGART HYTRULO can include hives.

- **Infusion or injection-related reactions.**

VYVGART can cause infusion-related reactions. VYVGART HYTRULO can cause infusion or injection-related reactions. These reactions can happen during or shortly after your VYVGART infusion or VYVGART HYTRULO injection. Tell your healthcare provider if you have any of the following symptoms of an infusion or injection-related reaction:

- high blood pressure
- chills
- shivering
- chest, stomach, or back pain

### The most common side effects of VYVGART or VYVGART HYTRULO include respiratory tract infection, headache, and urinary tract infection. An additional common side effect with VYVGART HYTRULO includes injection site reactions.

These are not all the possible side effects of VYVGART or VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART HYTRULO® (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines used to treat adults with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive. It is not known if VYVGART or VYVGART HYTRULO is safe and effective in children.

### Please see full Prescribing and Patient Information for VYVGART HYTRULO for subcutaneous injection and full Prescribing Information for VYVGART for IV infusion.

### Dosage forms and strengths:

VYVGART Hytrulo is available as a single-dose subcutaneous injection containing: 200 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per prefilled syringe, or 180 mg/mL of efgartigimod alfa and 2,000 U/mL of hyaluronidase per vial. VYVGART is available as a single-dose injection for intravenous use containing 400 mg/20 mL of efgartigimod alfa-fcab per vial.